

TU-539, RCNS-98-05
May 1998

Late-Time Entropy Production and Relic Abundances of Neutralinos

Takaaki Nagano and Masahiro Yamaguchi

*Department of Physics, Tohoku University,
Sendai 980-8578, Japan*

Abstract

Many models possess unwanted relics, which should be diluted by entropy production just before the big-bang nucleosynthesis. A field responsible for the entropy production may produce stable weakly interacting massive particles, if kinematically accessible. We compute their relic abundances by integrating out coupled equations numerically. Applying our results to supersymmetric standard models, we argue that the neutralino lightest superparticles will overclose the universe in most of the parameter space if the reheat temperature is of the order of 10 MeV.

Many models beyond the standard model predict unwanted relics, which should be diluted by entropy production such as inflation and subsequent reheating. Among other things, weakly interacting massive scalar fields will be the most problematic, as they can store their energy density in the form of coherent oscillation, which may not be diluted by inflation because the coherent mode during the inflationary epoch is generally displaced from its value in the true vacuum. An example is a problem associated with moduli fields of string theory, named the cosmological moduli problem [1]. In string theory coupling constants are determined dynamically by vacuum expectation values of the moduli (and dilaton) fields. These fields generically acquire masses comparable to the gravitino mass, which should be in the TeV range in gravity mediated supersymmetry breaking. Damped coherent oscillation and subsequent decay of the fields would spoil the standard big-bang nucleosynthesis (BBN). Late-time entropy production before the BBN should operate to dilute the energy density of the problematic scalar fields. Indeed thermal inflation was proposed [2, 3], for which reheat temperature should be much lower than the electroweak scale for a successful dilution.

Another example of the problematic scalar fields is an invisible axion with decay constant larger than 10^{13} GeV, whose coherent oscillation would exceed the critical density of the universe [4]. Entropy production may be able to dilute the energy density of the axion field [5]. Since the oscillation of the axion field starts around the QCD phase transition, the reheat temperature at the entropy production should be lower than 100 MeV. Interestingly it has been argued that strongly coupled $E_8 \times E_8$ heterotic string theory (M-theory) possesses such an axion candidate with the decay constant $\sim 10^{16}$ GeV, provided that non-perturbative effects to generate the superpotential for the axion field are suppressed [6, 7]. To dilute the energy density of the M-theory axion to a harmless level, the reheat temperature of the entropy production should be at its lowest value allowed by the BBN, 1–10 MeV.

The requisite entropy production is provided by decays of massive unstable particles. The decays, however, may produce too many stable particles as well if the production is kinematically allowed. In a supersymmetric theory, the lightest superparticle (LSP), which is likely to be a neutralino, is stable under the assumption of R -parity conservation. Indeed it was argued that the neutralinos produced by the decays of the massive particles tend to overclose the universe, if the reheat temperature is much lower than 1 GeV [8, 9].

Motivated by the aforementioned demands for the late-time entropy pro-

duction mechanism, we shall reexamine the problem of the relic abundances of the neutralinos which are produced by the decays of massive unstable particles. Given an annihilation cross section of the neutralinos, we will solve coupled equations numerically to evaluate the relic abundances, taking into account effects of exponential decays of the unstable particles, whereas an approximation of simultaneous decay was made in the previous works [8, 9]. It turns out that results obtained in our numerical calculation agree with the previous estimates up to a numerical factor of order unity and hence we confirm the validity of arguments in Refs. [8, 9]. Note that our computation should be applicable to any stable weakly-interacting-massive-particles. We will then argue that the stable LSP suffers from the overclosure problem in most of the parameter space if the reheat temperature is of the order of 10 MeV or less. We will also briefly mention possible implications of our analysis when there remains the M-theory axion in the massless spectrum of the heterotic M-theory.

We consider the situation that massive unstable particles ϕ of the decay width Γ come to dominate the energy density of the universe, then decay to neutralinos χ as well as to radiation with branching ratios B and $1 - B$, respectively. By assuming two-body decay, the energy density of χ just after the production is $m_\phi/2$. They subsequently lose their energy, by the scattering with thermal bath, down to $\simeq m_\chi (\gg T)$. Assuming this energy transportation occur rapidly, we define the effective branching ratio $B_{\text{eff}} \equiv B \frac{m_\chi}{m_\phi/2}$.

Time evolution of cosmic scale factor R and energy densities ρ_ϕ , ρ_R , ρ_χ of ϕ , radiation, and χ , respectively, are described by the following Friedmann equation and (integrated) Boltzmann equations:

$$H^2 = \left(\frac{\dot{R}}{R} \right)^2 = \frac{1}{3M^2} (\rho_\phi + \rho_R + \rho_\chi) , \quad (1)$$

$$\frac{d}{dt} (R^3 \rho_\phi) = -\Gamma (R^3 \rho_\phi) , \quad (2)$$

$$\frac{d}{dt} (R^3 \rho_R) + \frac{\rho_R}{3} \frac{dR^3}{dt} = (1 - B_{\text{eff}}) \Gamma (R^3 \rho_\phi) + \frac{\langle \sigma v \rangle}{m_\chi} (R^3 \rho_\chi)^2 \frac{1}{R^3} , \quad (3)$$

$$\frac{d}{dt} (R^3 \rho_\chi) = B_{\text{eff}} \Gamma (R^3 \rho_\phi) - \frac{\langle \sigma v \rangle}{m_\chi} (R^3 \rho_\chi)^2 \frac{1}{R^3} , \quad (4)$$

where M is the reduced Planck mass: it is related to the Newton's constant as $G = 1/m_{\text{Pl}}^2 = 1/(8\pi M^2)$, and $\langle \sigma v \rangle$ is an ensemble-averaged annihilation

cross section of neutralinos times their relative velocity. We treat $\langle\sigma v\rangle$ as a constant, which is indeed the case when the S -wave is dominant. When a higher partial wave dominates, on the other hand, $\langle\sigma v\rangle$ should be interpreted as a sort of weighted average in the time interval around when the annihilation process is important. In the above equations, we did not include pair production of the neutralinos, which is not effective at low temperatures. Note that the right-hand side of the Boltzmann equation for ρ_R is consistent with the energy conservation within comoving volume for the case $\dot{R} = 0$. Following [10] we introduce the dimensionless variables:

$$x \equiv \Gamma t, \quad z \equiv R/R_i, \quad (5)$$

$$f_\phi \equiv \rho_\phi/\rho_{\phi i}, \quad f_R \equiv \rho_R/\rho_{\phi i}, \quad f_\chi \equiv \rho_\chi/\rho_{\phi i}, \quad (6)$$

$$F_\phi \equiv z^3 f_\phi, \quad F_R \equiv z^4 f_R, \quad F_\chi \equiv z^3 f_\chi, \quad (7)$$

$$x_H \equiv \Gamma H_i^{-1} = \Gamma \left[\frac{\rho_{\phi i}}{3M^2} (1 + f_{Ri} + f_{\chi i}) \right]^{-1/2} \quad (8)$$

$$\rho_{\chi 0} \equiv \frac{m_\chi \Gamma}{\langle\sigma v\rangle}, \quad f_{\chi 0} \equiv \rho_{\chi 0}/\rho_{\phi i}. \quad (9)$$

Here subscripts i mean their initial values. By using these variables, the coupled differential equations above can be written as follows:

$$\frac{z'}{z} = x_H^{-1} (1 + f_{Ri} + f_{\chi i})^{-1/2} (z^{-3} F_\phi + z^{-4} F_R + z^{-3} F_\chi)^{1/2}, \quad (10)$$

$$F'_\phi = -F_\phi, \quad (11)$$

$$F'_R = (1 - B_{\text{eff}}) z F_\phi + f_{\chi 0}^{-1} z^{-2} F_\chi^2, \quad (12)$$

$$F'_\chi = B_{\text{eff}} F_\phi - f_{\chi 0}^{-1} z^{-3} F_\chi^2, \quad (13)$$

where prime denotes d/dx . We impose the following initial conditions at $x = x_i$,

$$z_i = 1, \quad F_{\phi i} = 1, \quad (14)$$

$$F_{Ri} = \frac{\rho_{Ri}}{\rho_{\phi i}} (\ll 1), \quad F_{\chi i} = \frac{\rho_{\chi i}}{\rho_{\phi i}} (\ll 1). \quad (15)$$

There is no difficulty in solving the Boltzmann equation for F_ϕ :

$$F_\phi = e^{-(x-x_i)}. \quad (16)$$

By assuming $\rho \propto R^{-3}$ for $x_i \lesssim x \lesssim 1$,

$$z^{3/2} = \frac{3}{2} \frac{x}{x_H} \quad \text{and} \quad \Gamma H^{-1} = \frac{3}{2} x \quad (17)$$

for $x_i \ll x \lesssim 1$. Our numerical calculation shows that $z|_{x=1}$ is about 10% larger than the estimate $(\frac{3}{2}x_H^{-1})^{2/3}$.

With an approximation of the simultaneous ϕ decay at $x = 1$, one obtains

$$\begin{aligned} F_R|_{x \gg 1} &= z|_{x=1} \cdot F_\phi|_{x \ll 1} \\ &\simeq \left(\frac{3}{2}x_H^{-1}\right)^{2/3} \\ &= 1.20 \times (1.09 x_H^{-2/3}) . \end{aligned} \quad (18)$$

Numerical calculation [10] for the coupled equations shows, on the other hand,

$$\begin{aligned} F_R|_{x \gg 1} &= \int_0^\infty z e^{-x} dx \\ &= 1.09 x_H^{-2/3} , \end{aligned} \quad (19)$$

for $F_{Ri} < 10^{-3}$ and $B_{\text{eff}} < 10^{-3}$. We confirm that the formula above is correct also for larger F_{Ri} and/or B_{eff} with at most about 10% error.

Our main concern in this paper is F_χ for $x \gg 1$, to which ρ_χ is proportional. A crude behavior of F_χ can be seen as follows. If $f_{\chi 0}^{-1} = \frac{\langle \sigma v \rangle}{m_\chi \Gamma} \rho_{\phi i}$ is negligibly small, or quantitatively if $B_{\text{eff}} x_H^2 f_{\chi 0}^{-1} < 1$ as we shall see shortly,

$$F_\chi = \begin{cases} B_{\text{eff}}(x - x_i) + F_{\chi i} & , \text{ for } x_i < x \lesssim 1 \\ B_{\text{eff}} & , \text{ for } x \gtrsim 1 . \end{cases} \quad (20)$$

That is, F_χ increases monotonically with x . On the other hand, if $f_{\chi 0}^{-1}$ is not so small the second term of the Boltzmann equation for F_χ cannot be neglected when F_χ becomes large:

$$F'_\chi|_{x=1} \simeq B_{\text{eff}} e^{-1} [1 - 1.2 B_{\text{eff}} x_H^2 f_{\chi 0}^{-1}] . \quad (21)$$

Thus when $B_{\text{eff}} x_H^2 f_{\chi 0}^{-1} > 1$, F_χ increases until $x \sim 1$, but decreases thereafter to reach $F_\chi|_{x \gg 1} < B_{\text{eff}}$. Note that

$$\begin{aligned} B_{\text{eff}} x_H^2 f_{\chi 0}^{-1} &= 6 M^2 B \Gamma \frac{\langle \sigma v \rangle}{m_\phi} \\ &= 1.17 \left(\frac{\langle \sigma v \rangle}{10^{-10} \text{GeV}^{-2}} \right) \left(\frac{m_\phi}{10^3 \text{GeV}} \right)^{-1} \left(\frac{B}{0.5} \right) \left(\frac{\Gamma^{-1}}{1 \text{sec}} \right)^{-1} . \end{aligned} \quad (22)$$

Fig. 1 shows the evolution of F_χ , obtained by integrating out the coupled differential equations numerically. It shows it increases till $x = 2$ or so, then decreases to reach a constant value. In the same figure, we also show the evolutions of the other quantities, z , F_ϕ and F_R .

Let us now compare our numerical results with a naive estimate given in [8]. This naive estimation of $F_\chi|_{x \gg 1}$ for the case where the annihilation process becomes effective was obtained by equating the annihilation rate $\langle \sigma v \rangle \rho_\chi / m_\chi$ to the expansion rate H at the decay epoch ($x = 1$) of ϕ , with the approximation of the simultaneous decay. One finds

$$\rho_\chi|_{x=1} = \left. \frac{m_\chi H}{\langle \sigma v \rangle} \right|_{x=1} = \frac{2}{3} \frac{m_\chi \Gamma}{\langle \sigma v \rangle} , \quad (23)$$

or,

$$F_\chi|_{x \gg 1} = F_\chi|_{x=1} = \frac{3}{2} x_H^{-2} f_{\chi 0} . \quad (24)$$

On the other hand, we find the following fit for our numerical results (see Fig. 2):

$$F_\chi|_{x \gg 1} = 1.52 (B_{\text{eff}} x_H^2 f_{\chi 0}^{-1})^{-0.919} B_{\text{eff}} \quad (25)$$

for $10 < B_{\text{eff}} x_H^2 f_{\chi 0}^{-1} < 10^8$ within 15% error. Comparison of the two results shows that the naive estimate is correct up to a factor of order unity in a wide range of the parameters. As $B_{\text{eff}} x_H^2 f_{\chi 0}^{-1} \propto B \Gamma \frac{\langle \sigma v \rangle}{m_\phi}$ becomes larger, $F_\chi|_{x \gg 1}$ is more underestimated if one uses the naive estimation. For example, for $B_{\text{eff}} = 10^{-1}$ and $x_H^2 f_{\chi 0}^{-1} = 10^8$,

$$F_\chi|_{x \gg 1}^{\text{naive}} = 1.5 \times (10^8)^{-1} = 1.5 \times 10^{-8} , \quad (26)$$

while,

$$\begin{aligned} F_\chi|_{x \gg 1}^{\text{numerical}} &= 1.52 \times (10^{-1} \times 10^8)^{-0.919} \times 10^{-1} \\ &= 5.61 \times 10^{-8} \\ &= 3.7 \times F_\chi|_{x \gg 1}^{\text{naive}} . \end{aligned} \quad (27)$$

Now we turn to $\rho_\chi / s|_{x \gg 1}$ where s stands for entropy density. Using

$$\begin{aligned} \rho_R &= a T^4 , \\ s &= \frac{4}{3} \frac{\rho_R}{T} = \frac{4}{3} a^{1/4} \rho_R^{3/4} , \\ a &= \frac{\pi^2}{30} g_* \end{aligned} \quad (28)$$

with g_* being the effective relativistic degrees of freedom, we find

$$\frac{\rho_\chi}{s} = \frac{3}{4} \frac{\rho_{\phi i} F_\chi}{a^{1/4} F_R^{3/4}} = \frac{3^{5/4}}{4} \frac{F_\chi}{a^{1/4} (x_H^{2/3} F_R)^{3/4}} (M\Gamma)^{1/2} . \quad (29)$$

Substituting our numerical results Eqs. (19) and (25), we obtain the following fit

$$\begin{aligned} \left. \frac{\rho_\chi}{s} \right|_{x \gg 1} &= \frac{3^{5/4}}{4} (1.09)^{-3/4} \left(\frac{\pi^2}{30} g_* \right)^{-1/4} (M\Gamma)^{1/2} \\ &\quad \times 1.52 \left(6M^2 B \Gamma \frac{\langle \sigma v \rangle}{m_\phi} \right)^{-0.919} \times 2B \frac{m_\chi}{m_\phi} \\ &= 3.08 h^{-2} \times 10^4 \left. \frac{\rho_{\text{cr}}}{s} \right|_0 \\ &\quad \times \left(\frac{\langle \sigma v \rangle}{10^{-10} \text{GeV}^{-2}} \right)^{-0.919} \left(\frac{m_\chi}{10^2 \text{GeV}} \right) \left(\frac{\Gamma^{-1}}{1 \text{sec}} \right)^{0.419} \\ &\quad \times \left(\frac{m_\phi}{10^3 \text{GeV}} \right)^{-0.081} \left(\frac{B}{0.5} \right)^{0.081} \left(\frac{g_*}{10.75} \right)^{-0.46} , \end{aligned} \quad (30)$$

where $\rho_{\text{cr}}/s|_0 = 3.64 h^2 \times 10^{-9} \text{GeV}$ is the ratio of the critical density ρ_{cr} to the entropy density s at present. This formula reproduces values of $\rho_\chi/s|_{x \gg 1}$ calculated numerically within 15% error for $10 < B_{\text{eff}} x_H^2 f_{\chi 0}^{-1} < 10^8$. Alternatively, the annihilation cross section is related to the relic abundance as

$$\begin{aligned} \langle \sigma v \rangle &= 0.767 h^{-2.18} \times 10^{-5} \text{GeV}^{-2} \\ &\quad \times \left(\frac{\rho_\chi/s|_{x \gg 1}}{\rho_{\text{cr}}/s|_0} \right)^{-1.09} \left(\frac{m_\chi}{10^2 \text{GeV}} \right)^{1.09} \left(\frac{\Gamma^{-1}}{1 \text{sec}} \right)^{0.456} \\ &\quad \times \left(\frac{m_\phi}{10^3 \text{GeV}} \right)^{-0.0881} \left(\frac{B}{0.5} \right)^{0.0881} \left(\frac{g_*}{10.75} \right)^{-0.501} . \end{aligned} \quad (31)$$

One finds that Eq. (31) agrees with the previous estimate in Ref. [8], up to a factor of order unity, if one recalls the temperature when ϕ decays is

$$T \simeq g_*^{-1/4} \sqrt{M\Gamma} \sim \left(\frac{\Gamma^{-1}}{1 \text{sec}} \right)^{-1/2} \text{MeV} , \quad (32)$$

which may be referred to as the reheat temperature.

The ensemble-averaged annihilation cross section (times relative velocity) $\langle\sigma v\rangle$ can be, in general, expanded in the averaged velocity squared $\langle v^2\rangle$ of the LSP:

$$\langle\sigma v\rangle = a + b\langle v^2\rangle + \cdots . \quad (33)$$

The coefficients a , b , *etc.* strongly depend on the composition of the LSP as well as superparticle mass spectra [11]. Sometimes the coefficient a of the S -wave becomes very small due to the Majorana nature of the neutralino. This is indeed the case when the LSP is the bino-dominant neutralino, in which a is proportional to the mass squared of the fermion in the final state. Then, in order to evaluate $\langle\sigma v\rangle$ and subsequently $\Omega_\chi h^2$, we have to know $\langle v^2\rangle$ precisely. The authors of Ref. [9] found that the bino-dominant LSP is likely to reach kinetic equilibrium and thus $\langle v^2\rangle \simeq 3T/m_\chi$ when the reheate temperature T is somewhat high, $T \gtrsim$ a few $\times 10$ MeV. Using this value of $\langle v^2\rangle$, they concluded that the LSP would overclose the universe as far as the reheate temperature is lower than a few hundred MeV. It is not clear, however, whether the LSP is in the equilibrium or not for other sets of parameters. Without detailed knowledge of the velocity distribution, there are still some conclusions we can draw in general. For instance, in view of the fact that the annihilation cross section is generally bounded from above as

$$\langle\sigma v\rangle \lesssim \frac{\pi\alpha_2^2}{m_\chi^2} \sim \frac{10^{-2}}{m_\chi^2} \quad (34)$$

we may conclude that the LSP with the mass at a few tens GeV or more does not annihilate sufficiently enough to survive the overclosure limit, if the reheate temperature is relatively low, say $T \lesssim \mathcal{O}(10)$ MeV, or $\Gamma^{-1} \gtrsim \mathcal{O}(10^{-4})$ sec, which seems to be required to make the M-theory axion viable. Note that the bound (34) does not apply when s -channel resonances (by Z boson or Higgs bosons) occur. To discuss the relic abundance in this case, it seems that we need to evaluate the velocity distribution of the LSP, which is beyond the scope of the present paper.

Finally we would like to discuss implications of our analysis to the M-theory if it turns out that the axion with large decay constant indeed arises. One needs to invoke an entropy production mechanism to dilute the energy density of the axion's coherent oscillation, unless its initial amplitude can be tuned to be very small. As we argued, the entropy production would produce too many neutralinos, if kinematically allowed, which would not annihilate efficiently. As a result they would overclose the universe, if they remain today.

This suggests that R-parity should be violated, making the neutralino-LSP unstable. Of course there are many loopholes to this argument, and we would not insist that the R-parity must be broken in the heterotic M-theory. Rather we regard the R-parity violation as an interesting option to which more attention should be paid.

Acknowledgments

This work was supported in part by the Grant-in-Aid for Scientific Research from the Ministry of Education, Science and Culture of Japan No. 09640333.

References

- [1] B. de Carlos, J.A. Casas, F. Quevedo and E. Roulet, Phys. Lett. **B318** (1993) 447;
T. Banks, D.B. Kaplan and A.E. Nelson, Phys. Rev. **D49** (1994) 779.
- [2] D.H. Lyth and E.D. Stewart, Phys. Rev. **D53** (1996) 1984.
- [3] E.D. Stewart, M. Kawasaki and T. Yanagida, Phys. Rev. **D54** (1996) 6032.
- [4] J. Preskill, M. Wise and F. Wilczek, Phys. Lett. **B120** (1983) 127;
L. Abbott and P. Sikivie, Phys. Lett. **B120** (1983) 133;
M. Dine and W. Fischler, Phys. Lett. **B120** (1983) 137.
- [5] M. Dine and W. Fischler, in Ref. [4];
P.J. Steinhardt and M.S. Turner, Phys. Lett. **B129** (1983) 51;
K. Yamamoto, Phys. Lett. **B161** (1985) 289;
J.E. Kim, Phys. Rev. Lett. **67** (1991) 3465;
M. Kawasaki, T. Moroi and T. Yanagida, Phys. Lett. **B383** (1996) 313;
M. Hashimoto, K.-I. Izawa, M. Yamaguchi and T. Yanagida, preprint hep-ph/9803263.
- [6] T. Banks and M. Dine, Nucl. Phys. **B479** (1996) 173; Nucl. Phys. **B505** (1997) 445.
- [7] K. Choi, Phys. Rev. **D56** (1997) 6588.

- [8] T. Moroi, M. Yamaguchi and T. Yanagida, Phys. Lett. **B342** (1995) 105.
- [9] M. Kawasaki, T. Moroi and T. Yanagida, Phys. Lett. **B370** (1996) 52.
- [10] R.J. Scherrer and M.S. Turner, Phys. Rev. **D31** (1984) 681.
- [11] M. Drees and M.M. Nojiri, Phys. Rev. **D47** (1993) 376.

Figure captions

Figure 1 Time evolution ($x \equiv \Gamma t$) of $z \equiv R/R_i$, $F_\phi \equiv z^3 f_\phi$, $F_R \equiv z^4 f_R$, $F_\chi \equiv z^3 f_\chi$, and ρ_χ/s , where $f_* \equiv \rho_*/\rho_{\phi i}$ ($*$ = ϕ , R , χ) and Γ is the decay width of ϕ , for the case when $B_{\text{eff}} \equiv 2m_\chi/m_\phi = 10^{-1}$, $x_H \equiv \Gamma H_i^{-1} = 10^{-2}$, $f_{\chi 0} \equiv m_\chi \Gamma / (\langle \sigma v \rangle \rho_{\phi 0}) = 10^{-9}$, $\Gamma^{-1} = 1\text{sec}$, and $g_* = 10.75$. We can see that entropy $S \propto F_R^{3/4}$ within comoving volume increases during $0.01 \lesssim x \lesssim 2$. We can also see F_χ increases until $x \simeq 2$ but decreases thereafter to reach certain constant value. Likewise, ρ_χ/s decreases for $x \gtrsim 2$ to reach certain asymptotic value.

Figure 2 Asymptotic values of F_χ and ρ_χ/s as a function of $f_{\chi 0}^{-1}$, for the case $B_{\text{eff}} = 10^{-1}$, $x_H = 10^{-2}$, $\Gamma^{-1} = 1\text{sec}$, and $g_* = 10.75$. Qualitative behavior of $F_\chi|_{x \gg 1}$ and $\rho_\chi/s|_{x \gg 1}$ changes depending on if $B_{\text{eff}} x_H^2 f_{\chi 0}^{-1}$ is greater than 1. We also show naive estimations, $F_\chi|_{x \gg 1}^{\text{naive}} = (3/2) x_H^{-2} f_{\chi 0}$ and $\rho_\chi/s|_{x \gg 1}^{\text{naive}} = 2^{-3/2} 3^{3/4} a^{-1/4} \sqrt{M\Gamma} F_\chi|_{x \gg 1}^{\text{naive}}$, and linear fits for $10 < B_{\text{eff}} x_H^2 f_{\chi 0}^{-1} < 10^8$, for comparison. Linear fits are accurate within 15% error.

$$B_{\text{eff}} = 10^{-1}, \quad x_H = 10^{-2}, \quad f_{\chi 0} = 10^{-9}, \quad \Gamma = (1 \text{ sec})^{-1}, \quad g_* = 10.75$$




Figure 1

$$B_{\text{eff}} = 10^{-1}, x_H = 10^{-2}, \Gamma = (1 \text{ sec})^{-1}, g_* = 10.75$$




Figure 2